

Feltrin L, Bertelli M, McGaughey J, Morris WA, Piercey SJ, Kyser TK, Ansdell K, Galley A, Lesher CM, Weber D, Ferreira P, Quirt DH, Kotzer TG, 2016, HYPERCUBE: mining exploration data, a case study using the Millennium uranium deposit, Athabasca basin, Abstract, Geostats, Valencia, Italy, 1, 295-296

The Canada Mining Innovation Council (CMIC) Data Integration Technology Group, as part of a five-year commitment to research and innovation in mineral exploration (the Footprints Project), is applying predictive analytics to characterize quantitatively the footprints of well-known ore deposits in Canada. Data mining was used to evaluate meaningful associations in geological, geophysical, geochemical, mineralogical and petrophysical data at numerous sites and here we report initial results on a case-study, conducted on the Millennium uranium deposit (indicated resource of 75.9 million pounds U₃O₈ at grade of 2.39% U₃O₈, inferred resource of 29 million pounds U₃O₈ at grade of 3.19% U₃O₈). Market basket analysis was used to establish sets of “association rules” among 3D variables, including litho-geochemical, mineralogical, and geophysical variables, at various distances from the core to periphery at the Millennium deposit. The methodology implemented identifies “association rules” using BearingPoint, HyperCube® (HC) rule mining technology. Different ranking statistics were applied to select optimal rule-sets in the mineralized basement and in the overlying sandstone units of the Manitou Falls Formation. To evaluate HC results, systematic exploratory data analysis (EDA) was conducted to examine observed associations for all the variables considered. Results obtained suggest that HC correctly identifies the change in U associations due to compositional variations attributed to different mineralogy of the host-rock. Together, EDA and HC analyses show positive correlation of U with ²⁰⁶Pb/²⁰⁴Pb and ²⁰⁷Pb/²⁰⁴Pb ratios, Mo, and to a lesser extent V, in all analyzed stratigraphy: indicative of primary dispersion. Positive associations of U and HREE+Y are dominant in sandstones above the deposit, whereas at the fringes of the deposit, U is associated with Rb, Pb, and Th in the basement. Anticorrelations with U were recorded for major/trace elements in the basement (e.g., Mn, Si, Ca, Ba, Na, and Fe), reflecting the transition from altered clay mineral-dominated assemblages to unaltered rocks with plagioclase and K-feldspar. Anticorrelations with U in sandstone-cover for Ce, Th, Zr, and Hf reflect a transition from U phases within detrital minerals (e.g., zircon, monazite, apatite) to U in hydrothermal phases such as uraninite, APS, and other REE-hosting minerals. Furthermore, HC simple lift analysis indicates that radiogenic ²⁰⁶Pb/²⁰⁴Pb ratios higher than 30 are located in the MFd unit above the deposit: consistent with secondary dispersion of radiogenic Pb several hundreds of metres from the deposit.

NSERC-CMIC Mineral Exploration Footprints Project Contribution 090.