

***Lypaczewski P, Gaillard N, Perrouty S, Rivard B, Linnen RL, 2016, Vectoring towards mineralization using hyperspectral imaging at the Canadian Malartic gold deposit, Quebec, Canada, Thesis Presentation, U Laval, QC***

The Canadian Malartic gold deposit is mainly hosted in Archean metasediments, which are challenging to characterize by conventional core logging. We made use of a hyperspectral imaging system (Specim SisuROCK™) to acquire shortwave infrared (SWIR) and thermal infrared (TIR) reflectance spectra over 150 m of continuous drill core from the ore zone, as well as intervals of weakly altered, distal cores. The presence of quartz was determined from the TIR imagery, while the presence and mineral chemistry of white mica, biotite and chlorite was assessed with SWIR data. The drill core imagery enabled a correlation of estimated mineral compositions with downhole Au grades. The presence of phengitic white mica (>2206nm, <3.3 Al<sup>vi</sup> apfu) and Mg-rich biotite/chlorite (Mg# > 70) is indicative of mineralized intervals, whereas less altered, distal samples, are more muscovitic (<2202nm, >3.5Al<sup>vi</sup> apfu), with biotite/chlorite of an intermediate composition (Mg# 50-60). The SWIR analysis also included over 800 point measurements collected with a portable field spectrometer (Terraspec®) from outcrops in a 8 x 12 km region surrounding the deposit. This high density data allowed us to isolate the effect of metamorphism from that of hydrothermal alteration on mineral chemistry. These surface measurements revealed a multi-km hydrothermal alteration halo around the deposit that may be of value for vectoring towards mineralization in similar environments.

NSERC-CMIC Mineral Exploration Footprints Project Contribution 109.